UNIVERSITY HOSPITALS OF DERBY & BURTON NHS FOUNDATION TRUST DRUG MONOGRAPH FOR USE ON ADULT INTENSIVE CARE UNITS

ADDAVEN				
Indication	Patients who are nutritionally depleted & not receiving TPN or NG feed who require trace element supplementation. Refer to ICU Nutrition Guidelines.			
Dose	One ampoule ONCE daily			
Preparation	 Withdraw 10ml from a 100ml sodium chloride 0.9% bag Draw up 10ml ampoule of Addaven Add Addaven to sodium chloride 0.9% bag 			
Administration	Infuse over 2 hours via a peripheral or central line.			
Shelf-life	Use immediately			
Common Compatibility Issues	Addaven should be infused via a dedicated lumen Glucose 5% can be used as a diluent if required			
Additional information	Trace elements include zinc, selenium, copper and manganese.			
	DRUGS ADDED TO THIS INFUSION PATIENT A. Patlent (A. Number)			WARD ICH
Sample Label	DRUG Addaven ín 100ml sodíum chloríde 0.9%	AMOUNT 10ml	ADD BY	CHECKED BY
	DATE ADDED TIME ADDED	EXP. DATE EXP. TIME		BATCH No.
	DISCONTINUE IF CLOUDINESS OR PRECIPITATE DEVELOPS			

For review June 2025

Documentation Controls	
Development of Guideline:	Pharmacist – Critical Care & Theatres
Consultation with:	Pharmacy Department
Approved By:	ICU Sister's Meeting: June 2023 ICU Risk & Quality Meeting: August 2023
Deview Dates	Surgical Division
Review Date:	June 2025
Key contact:	Pharmacist – Critical Care & Theatres

References

Summary of Product Characteristics

<u>Microsoft Word - 3801863901315797365_spc-doc.doc (windows.net)</u> Accessed July 2022 Trissel, L.A. Handbook on Injectable Medicines 18th Edition.

https://www.medicinescomplete.com/mc/hid/current/index.htm Accessed July 2022. Brayfield A (ed), Martindale: The Complete Drug Reference. [online] London: Pharmaceutical Press <<u>http://www.medicinescomplete.com/</u>> (accessed July 2022). Fresnius Kabi, pers comm 04/05/2021

*** End of Monograph ***