

## Hypospadias - Paediatric Full Clinical Guideline - UHDB

Reference no.: WC/NP/44N

### 1. Introduction

Hypospadias is a congenital malformation of the male genitalia characterised by incomplete development of the urethra in which the opening is located on the ventrum rather than at the tip of the penis. Severity depends on the location of the urethral opening. In most cases, the opening is located on or close to the glans. Moderate or severe forms of hypospadias occur when the opening is situated more proximally on the penile shaft, at the base of the penis or, in rare cases, in the scrotum or perineum.

### 2. Aim and Purpose

To inform staff, within all areas, in the management of neonates and infants with hypospadias.

### 3. Classification

Hypospadias can be classified according to location (glandular, coronal, subcoronal, distal penile, mid shaft, proximal penile, peno-scrotal, scrotal or perineal) or according to severity i.e. mild (glandular or coronal), moderate (subcoronal or penile) and severe (peno-scrotal, scrotal or perineal).

### 4. Management

The aetiology of hypospadias remains unknown with environmental exposure in the form of endocrine disruptors being the most likely explanation. Appendix 1 is a general guideline on the management of newborns noticed to have hypospadias on the Postnatal Wards.

In cases of undescended testes together with hypospadias (proximal defect), chromosome analysis should be requested and on-call consultant to be informed. An urgent referral to Paediatric Urologist may be required.

Distal defects (e.g. hooded prepuce) can be referred to Urologist after reassessment in the clinic at 2-3 months of age.

Advise parents to avoid circumcision on religious grounds until reviewed by Paediatric Urologist.

## 5. Suggested key review criteria for monitoring and/or audit

Appropriateness of management/follow up according to the severity of hypospadias

Associated congenital/chromosomal anomalies

Timing of referral to Paediatric Urologist

Delayed diagnosis (beyond neonatal age group)

## 6. References

Abdullah NA, Pearce MS, Parker L, Wilkinson JR, Jaffray B and McNally RJQ. Birth prevalence of cryptorchidism and hypospadias in northern England, 1993–2000 Arch Dis Child. 2007;92:576-579

Ahmed SF, Dobbie R, Finlayson AR, Gilbert J, Youngson G, Chalmers J & Stone D. Prevalence of hypospadias and other genital anomalies among singleton births, 1988–1997, in Scotland Arch Dis Child Fetal Neonatal Ed. 2004;89:149-151

Leonard J. Paulozzi, J. David Erickson, and Richard J. Jackson  
Hypospadias Trends in Two US Surveillance Systems Pediatrics.  
1997;100:831-834

Baskin LS, Ebberts MB. Hypospadias: anatomy, etiology, and technique. *J Pediatr Surg* 2006;41 (3) :463–72

<b>Reference Number</b> <b>WC/NP/44N</b>	<b>Version:</b> V7		<b>Status</b> Final	
Version / Amendment History	Version	Date	Author	Reason
	V7	March 2024	Mr More	Review and update
<b>Intended Recipients:</b> All Paediatric Medical Staff, Nursing Staff and Midwives at Derby Hospital				
<b>Training and Dissemination:</b> Cascade the information via BU newsletter and address training				
<b>Development of Guideline:</b> Mr More Consultant Paediatric Urologist and Surgeon				
<b>In Consultation with:</b> All Paediatric Medical Staff and Nursing Staff plus Senior Nurse Managers and all midwives				
<b>Linked Documents:</b> 58: Policy Framework				
<b>Keywords:</b> Hypospadias				
<b>Business Unit Sign Off</b>			<b>Group:</b> Paediatric Guidelines Group <b>Date:</b> 05/06/2024	
<b>Divisional Sign Off</b>			<b>Group:</b> Women's and Children's Clinical Governance Group <b>Date:</b>	
<b>Date of Upload</b>			22 <sup>nd</sup> July 2024	
<b>Review Date</b>			March 2027	
<b>Contact for Review</b>			Dr B More	

## Appendix 1: Management of Hypospadias

