

## Dyspepsia - Full Clinical Guideline

Reference No: CG-T/2012/020

### **Aim**

To provide clinicians with guidance on managing patients with dyspepsia, aiming to assist with identification of patients who need to be investigated rapidly and those whom require lifestyle advice and reassurance about the benign nature of their condition. Whilst these guidelines are primarily aimed at the out-patient population, they can equally be applied to in-patients.

### **Purpose and Scope**

Dyspepsia can be defined as pain or discomfort that is attributable to the upper gastrointestinal (GI) tract and may be related to eating and drinking. The routine investigation of patients with dyspepsia is not always necessary.

**Any patient with the alarm symptoms listed below require urgent investigation with gastrointestinal endoscopy:**

- Dysphagia
- Age 55 and over with weight loss **and any of the following**:
  - upper abdominal pain
  - reflux
  - dyspepsia
  - heartburn
  - nausea or vomiting
  -

Consider non-urgent investigation with gastrointestinal endoscopy in:

- Age 55 or over **and one or more of the following**:
  - treatment resistant dyspepsia
  - upper abdominal pain and anaemia
  - raised platelet with any of the following;
    - nausea
    - vomiting
    - weight loss
    - reflux
    - dyspepsia
    - upper abdominal pain
  - nausea and vomiting **with**:
    - weight loss
    - reflux
    - dyspepsia
    - upper abdominal pain
- Haematemesis or Melaena

- In any age
  - Gastro-oesophageal symptoms that are non-responsive to treatment or unexplained
  - Suspected GORD considering anti-reflux surgery
  - *H pylori* that has not responded to second line eradication
  - Iron deficiency anaemia
  - Abnormal radiological investigation

Consider differential diagnoses such as gallbladder and pancreatic disease and ovarian cancer in women as they can present with dyspeptic symptoms.

### **Guideline**

- Consider urgent gastroscopy with criteria above
- Review medications for possible causes of dyspepsia e.g. calcium antagonists nitrates, theophyllines, bisphosphonates, NSAIDs, Corticosteroids
- Give lifestyle advice e.g. healthy eating, weight reduction and smoking cessation, avoid known precipitants e.g. smoking, alcohol, coffee, chocolate and fatty/spicy foods. Raising the head of the bed and having a main meal well before bedtime may help if symptoms are worse at night
- Test for *Helicobacter pylori* (using *H pylori* stool antigen test) and treat with eradication therapy (see BNF)
- Give a course of a proton pump inhibitor (PPI), if this relieves symptoms of heartburn it is diagnostic of gastro-oesophageal reflux disease (GORD). Follow-up with life style advice and suggest PPI prn
  - Step down medication so that the patient is on the lowest possible dose to manage their symptoms e.g. PPI – H2 receptor antagonist (H2RA) – Raft Preparation (antacid/alginate preparation) which can brought over the counter
  - Consider managing previously investigated patients without alarm symptoms according to previous endoscopic findings
  - Patients who have persistent symptoms may require double dose PPI plus H2RA at bedtime
  - Consider Laproscopic anti-reflux surgery
  - Patients without alarm symptoms (as listed above) and a response to PPI therapy do not require endoscopy but can be managed symptomatically as above

## **References**

Evidence Based Interventions: List 2 Guidance. (2020) Academy of Medical Royal Colleges. Available at: <https://www.aomrc.org.uk/evidence-based-interventions/> [accessed 06/05/22]

British National Formulary (2020) British National Formulary 2020 [online] Available at: <https://www.medicinescomplete.com/mc/bnf/current/> [accessed 28.10.20]

Delaney, B. et al (2008) Helicobacter pylori test and treat versus proton pump inhibitor in initial management of dyspepsia in primary care: multicentre randomised controlled trial (MRC-CBUE trial) BMJ, 336: 651-654

Derbyshire Joint Area Prescribing Committee (JAPC) 2015 (reviewed 2017) Management of Dyspepsia and Gastro-oesophageal Reflux Disease (GORD) [on line] Available at: [https://www.derbyshiremedicinesmanagement.nhs.uk/assets/clinical\\_GuidelinesFormulary](https://www.derbyshiremedicinesmanagement.nhs.uk/assets/clinical_GuidelinesFormulary) [accessed 05.02.18]

National Institute for Health and Clinical Excellence (2014) Gastro-oesophageal Reflux Disease and Dyspepsia in Adults: investigation and management [on line] Available at: <https://www.nice.org.uk/guidance/cg184>

## **Keywords**

Dyspepsia

Adults

GORD

Proton Pump Inhibitor

Helicobacter pylori

**Documentation Control:**

Development of Guideline:	Dr J Williams, Consultant Gastroenterologist Ann Stokes, Gastroenterology Nurse Specialist
Consultation with:	Prof A Goddard, Consultant Gastroenterologist Dr E Tucker, Consultant Gastroenterologist
Approved By:	Gastroenterology Sub Directorate Group 20/01/21 Endoscopy Users Group 08/6/21 Medical Division – 02/02/21
Review Date:	May 2024
Key Contact:	Ann Stokes – Dyspepsia Nurse Specialist Tel No. 01332 786754