

Neonatal Seizures

Advances in Mechanisms and Management

Hannah C. Glass, MDCM, MAS

KEYWORDS

- Brain injury • Developmental disability • Infant, newborn • Electroencephalography
- Epilepsy • Magnetic resonance imaging • Neurocritical care • Seizures

KEY POINTS

- Seizures occur in 1 to 5 per 1000 live births and are among the most common neurologic conditions managed by a neonatal neurocritical care service.
- The high rate of seizures in the neonatal period reflects age-specific developmental mechanisms that lead to relative excitability.
- Neonatal seizures are often caused by serious underlying brain injury such as hypoxia-ischemia, stroke, or hemorrhage.
- Clinical detection is unreliable; continuous video electroencephalogram is the gold standard for monitoring presence and burden of neonatal seizures.
- Seizures are refractory to first-line medications in approximately 50% of cases; expert opinion supports rapid treatment to abolish acute symptomatic seizures and early discontinuation of medication.

INTRODUCTION

Neonates are at especially high risk for seizures as compared to other age groups.¹ The high risk for seizures—and especially acute symptomatic seizures—is likely multifactorial and often caused due to the relative excitability of the developing neonatal brain as well as the high risk for brain injury due to global hypoxia-ischemia, stroke, and intracranial hemorrhage.² The estimated rate of seizures in term newborns is said to be approximately 1 to 5 per 1000 live births.^{3–5} However, population-based studies do not take into account the low diagnostic accuracy of diagnosis by clinical

Financial Disclosure: Nothing to disclose.

Disclosures: None.

Project Support: H.C. Glass is supported by the NINDS K23NS066137, the Pediatric Epilepsy Research Foundation, and the Neonatal Brain Research Institute.

Departments of Neurology and Pediatrics, University of California, 505 Parnassus Avenue, M793, Box 0114, San Francisco, CA 94143-0114, USA

E-mail address: Hannah.Glass@ucsf.edu

observation alone,^{6,7} and gold standard prolonged, continuous video electroencephalogram (cEEG) monitoring is not widely available enough to make population-based predictions; therefore the true incidence remains unknown.

The differential diagnosis for neonatal seizures is broad and includes structural, metabolic, and genetic causes (**Box 1**). Seizures that arise from an acute symptomatic cause, such as hypoxic-ischemic encephalopathy, transient metabolic disturbance, infection, stroke, or intracranial hemorrhage, are much more common than neonatal onset epilepsies, which may be due to malformation, prior injury, or genetic causes. Rare conditions such as inborn errors of metabolism, vitamin-responsive epilepsies, and neonatal epilepsy syndromes must be considered in the setting of refractory seizures.^{8,9}

Neonatal seizures carry a high risk for early death. Among survivors, motor and cognitive disabilities, as well as epilepsy are common.¹⁰ The outcome depends largely on the underlying disease process and severity of underlying brain injury. The impact of the seizures themselves is not known, although studies in animal models suggest that seizures can alter brain development, leading to deficits in learning, memory, and behavior.¹¹

Box 1

Cause of neonatal seizures

Differential Diagnosis of Acute Symptomatic Seizures

Global hypoxia-ischemia (hypoxic-ischemic encephalopathy)

Focal hypoxia-ischemia

Arterial stroke

Venous stroke

Intracranial hemorrhage

Intraventricular

Parenchymal

Subarachnoid

Subdural

Transient metabolic deficit

Hypoglycemia

Hypocalcemia and hypomagnesemia

Hyponatremia

Acute infection

Differential Diagnosis of Neonatal Onset Epilepsy

Brain malformation

Intrauterine injury or congenital infection

Inborn error of metabolism and vitamin-responsive epilepsies

Neonatal Onset Epilepsy Syndromes

Benign familial neonatal seizures (eg., KCNQ2, KCNQ3)

Neonatal epileptic encephalopathies

Early myoclonic epilepsy

Early infantile epileptic encephalopathy (Ohtahara syndrome)

PATOPHYSIOLOGY

There are several, age-specific factors that are particular to the neonatal brain that lead to enhanced excitability and seizure generation, poor response to conventional medications, and adverse impact on brain development.¹²

Enhanced Excitability of the Neonatal Brain

There are numerous mechanisms that render the immature brain hyperexcitable as compared to the adult brain.^{12,13} First, the neonatal period is a time of physiologic, use-dependent synaptogenesis, and both synapse and dendritic spine density are at their peak.^{14,15} Second, glutamatergic neurons—the primary excitatory mechanism of both the developing and adult brain—are overabundant, and their receptors are configured with subunits that allow relative hyperexcitability.^{16,17} Third, gamma-aminobutyric acid (GABA)—the primary inhibitory mechanism of the adult brain—can exert a paradoxical excitatory action in the developing brain due to the preponderance of the NKCC1 and delayed expression of the KCC2 chloride cotransporters, which lead to a high intracellular chloride concentration and *depolarization* in response to GABAergic agents.^{18–20}

Anticonvulsants and the Developing Brain

Immature development of the excitatory and inhibitory neurotransmitter systems leads to a lack of good targets for conventional antiseizure medications, which makes neonatal seizures particularly difficult to treat. The immature brain may be resistant to medications that act as GABA agonists, not only as a result of the paradoxical chloride gradient as discussed earlier but also due to overall lower receptor expression and an immature subunit composition that is less sensitive to benzodiazepines than the adult brain.^{12,13}

Seizures and Early Brain Development

Although early work with animal models demonstrated that the developing brain is more resistant to seizure-induced necrosis than the adult brain, more recent work has shown that early-life seizures can affect the developing brain nonetheless by altering neuronal circuitry, which can result in impaired learning and memory and enhanced susceptibility to epilepsy later in life.¹¹ Animal models of early-life seizures display developmental alterations that can include reduced density of dendritic spines in hippocampal pyramidal neurons; decreased neurogenesis; delayed neuronal loss; and changes in hippocampal plasticity such as decreased capacity for long-term potentiation, reduced susceptibility to kindling, and enhanced paired-pulse inhibition.^{21–24} Human studies in children with hypoxic-ischemic injury show an independent association between seizures and impaired brain metabolism, as well as poor long-term neurodevelopmental outcome.^{25,26}

MANAGEMENT GOALS

The overall management goal for neonatal seizures is to quickly and accurately identify, and abolish electrographic seizures, while determining the most likely underlying cause. Neonatal seizures are often the first sign of neurologic dysfunction and are frequently an indication of serious underlying brain injury.^{27,28} Therefore, a suspicion of seizures in a newborn should be treated as a neurologic emergency, and prompt rapid and thorough evaluation to identify the cause, as well as emergent medical management to abolish seizures should be performed while preventing secondary injury by maintaining physiologic temperature, glucose, oxygenation, ventilation, and blood pressure.

Seizure Detection and Monitoring

Common methods for identifying neonatal seizures are outlined in **Table 1**. Clinical evaluation of seizures is approximately 50% accurate for events detected at the bedside. Furthermore, clinical detection requires constant observation by the bedside staff and even so will fail to detect seizures with no or very subtle clinical correlate (eg, eye deviation or subtle clonic movements that are covered by the infant's blanket). Subclinical seizures account for most seizures in neonates, especially in the setting of severe brain injury, and in children who have received seizure medications.^{7,29-31}

Recent guidelines from the American Clinical Neurophysiology Society set the standard for neurophysiology monitoring in neonates.³² cEEG, with electrodes placed according to the international 10–20 system, modified for neonates, is the gold standard for monitoring.³³⁻³⁵ Barriers to implementing this technology include the need for specialized training for the application and interpretation of the recording, as well as variable access to equipment, and high cost. Once initiated, cEEG should be maintained until electrographic seizures have resolved for at least 24 hours or 3 to 4 clinical events have been captured and determined not to be seizures.³²

Amplitude-integrated electroencephalography (aEEG; a simplified bedside neurophysiology tool that can be applied and interpreted by neonatologists, nurses, or other Intensive Care Nursery bedside staff) is used to supplement or even replace cEEG in a growing number of centers. Because aEEG uses a limited number of channels to record EEG signal that is heavily processed (filtered, rectified, and displayed on a semilogarithmic amplitude and time-compressed scale), there are several limitations to this technology that must be taken into account when it is used for management of neonatal seizures (see **Table 1**). Machines that allow for concurrent monitoring and display of aEEG (at the bedside) and cEEG (in the neurophysiology laboratory or remotely) using the same hardware have been suggested as a way to optimize use of both technologies, such that both the bedside team and the neurologist can be readily involved in the rapid, real-time management of electrographic seizures (**Fig. 1**).³⁶

Automated seizure detection—with an alarm to alert the bedside practitioner in the case of suspected seizures—is an attractive option that may offer the most practical solution for wide-scale implementation of seizure detection and management. However, there have been several challenges that limit the development of automated detection algorithms, including highly variable nature of neonatal seizure patterns,

Table 1
Diagnosis of neonatal seizures

Conventional cEEG	<ul style="list-style-type: none"> • Gold standard for seizure detection • Recommended by the American Clinical Neurophysiology Society for monitoring neonates with paroxysmal events and/or at high risk for seizures³²
aEEG	<ul style="list-style-type: none"> • Lower sensitivity and specificity than cEEG³⁶ • Hundred percent sensitivity for status epilepticus³⁷ • Lowest sensitivity for seizures that are brief, focal, and distal from recording electrodes^{38,39} • Raw EEG tracing helps to distinguish artifact from seizure⁴⁰ • Experienced readers perform better than nonexperts⁴¹
Clinical evaluation	<ul style="list-style-type: none"> • Accuracy approximately 50%⁶ • Will not identify most seizures (ie, subclinical or nonconvulsive seizures)^{7,29-31}




Fig. 1. aEEG (left) and EEG (right) from a term male with multiple intracranial hemorrhages and seizures that were refractory to phenobarbital, 40 mg/kg, and fosphenytoin, 30 mg/kg, and abated after 60 mg/kg levetiracetam. aEEG and EEG are recorded from a single machine at the bedside. Data are displayed differently for different users: the bedside team sees the aEEG display at left, which shows long-term trends and allows a quick review of suspicious segments of EEG. The neurophysiologist can confirm the seizures through the review of conventional, neonatal montage EEG.

high frequency of potential artifacts in the intensive care nursery, and uncertainty regarding the gold standard against which algorithms are tested given the limited information regarding inter-rater reliability among human expert readers.⁴² Newer algorithms that use machine learning, as well as temporal and spatial weighting hold promise.^{43–46}

Diagnostic Evaluation

Initial evaluation of a neonate with suspected seizures should also focus on rapid identification of the cause. Emergent evaluation of serum glucose and risk factors for infection is an important first step, because hypoglycemia and bacterial meningitis can lead to permanent injury if left untreated.⁸ Additional bedside evaluations must include measurement and treatment of electrolyte disturbance. Comprehensive history and physical examination are important tools to assess for risk factors and signs of both common and rare causes of neonatal seizures. Further evaluation, including genetic testing, serum amino acids, ammonia, lactate, very-long-chain fatty acids, urine organic acids and sulfites, and cerebrospinal fluid studies for glucose, glycine, lactate, and neurotransmitters, as well as additional testing for inborn errors of metabolism may be warranted on a case-by-case basis, especially in the setting of medically refractory seizures of unknown cause or a burst-suppression pattern on EEG in a neonate without brain injury.

Detailed neuroimaging using magnetic resonance (MR) is essential to identify underlying injury or developmental abnormalities and to help clinicians and the family to better understand the prognosis.⁴⁷ Cranial ultrasound, which is readily available at the bedside in most units, is important for rapid initial assessment of a sick neonate to identify large space occupying lesions, such as hemorrhage, arteriovenous malformations, or hydrocephalus, but is insensitive for global and focal hypoxic-ischemic injury, especially in the days after the ictus. Computed tomography exposes the infant to ionizing radiation and provides inferior resolution to MRI in most settings, and so should be avoided.

PHARMACOLOGIC STRATEGIES

There are no evidence-based guidelines for the pharmacologic management of neonatal seizures.^{48,49} Expert opinion supports use of pharmacologic treatments with a goal of abolishing electrographic seizures, even those without clinical correlate.⁴² However, evidence is lacking regarding the relative benefit versus potential harm of anticonvulsants used to treat seizures in neonates, many of which can lead to neuronal apoptosis in animal models.⁵⁰

Because seizures are refractory to initial doses of medication in approximately 50% of cases,⁵¹ frequent reevaluation of cEEG and bedside monitoring is essential to accurately identify and treat ongoing seizures in real time. Although data are lacking regarding optimal treatment paradigms for neonatal seizures, experts advocate *rapid* administration of an adequate loading dose of medication because acute symptomatic seizure burden is highest at the onset,⁵² and patients with fewer seizures are easier to treat.⁵¹ Similarly, experts advocate treatment of both clinical *and* subclinical seizures given similar pathophysiology, and the only difference between the 2 may be slight anatomic differences in their cortical distribution.⁴² Use of algorithms or guidelines to direct the treatment of neonatal seizures has gained favor, given evidence that treatment guidelines can improve outcomes in other settings.⁵³ As discussed later in this article, the optimal medication for seizure therapy in neonates is not known, and so guidelines should focus on an institution-specific, consensus-based protocol with

input and acceptance by both neonatology and neurology services to help prevent unnecessary delays in treatment that may result from discussions over medication choice in the setting of an actively seizing neonate.

The optimal duration of pharmacologic therapy for acute symptomatic seizures is not known. Treatment practices are variable in spite of good evidence that there is no harmful effect of early discontinuation of seizure therapy and no difference in seizure recurrence risk among neonates who are maintained on therapy versus those whose medication is maintained until several months of age (**Box 2**).^{54–56}

International survey data support the use of phenobarbital as the first-line medication based on expert consensus (**Table 2**).^{57–59} A single randomized, controlled trial found that phenobarbital and phenytoin were equally efficacious as first-line agents for seizure cessation among term infants with seizures.⁵¹ However, seizure control (defined by the study parameters as an 80% reduction in the severity of seizures) was achieved in fewer than half of the infants.⁵¹ This result is supported by newer studies, which demonstrate that up to 50% of neonatal seizures are refractory to first-line medications and an additional 30% fail second-line therapy.⁶⁰

Levetiracetam is gaining increasing support, in spite of limited efficacy data^{61,62}; this is likely due to the ready availability of an intravenous formulation in the United States, as well as a favorable safety and tolerability profile among children and adults.⁶³ In contrast to older agents such as phenobarbital and phenytoin, levetiracetam does not appear to enhance neuronal apoptosis in animal models^{64,65} and may in fact have neuroprotective and antiepileptogenic effects.^{66,67} The optimal neonatal dosage of levetiracetam is not yet known: reported doses range from 5 to 60 mg/kg/d.⁶⁸ However, the high volume of distribution and rapid clearance in neonates may necessitate a higher loading dose and more frequent dosing to maintain serum concentrations in the range used for adults and children.^{69,70} Published studies of the clinical efficacy of levetiracetam that report seizure reduction or resolution in 35% to 80% are limited by lack of standardized dosing, limited EEG monitoring, no placebo comparison, and/or variable timing and definition for determining the outcome.^{71–73}

Common agents for refractory seizures include midazolam infusion, which may be effective for neonatal status epilepticus, and lidocaine, which is widely used for refractory neonatal seizures in Europe.^{74–81} Topiramate is an antiseizure medication that has multiple mechanisms of anticonvulsant action and is an interesting option for acute symptomatic neonatal seizures because it appears to have neuroprotective effects in animal models of seizures and brain injury.^{82,83} A recently developed intravenous preparation of topiramate that is well tolerated in adult volunteers and has equivalent bioavailability to the oral formulation holds promise for use in neonates.^{84,85}

Bumetanide is a loop diuretic that has been proposed as an adjunct to GABAergic drugs like phenobarbital to help overcome the depolarizing action of immature neurons to GABA agonists. The mechanism of action is presumed to be through reduction in intracellular chloride concentrations, thus rendering the normally excitatory response of immature cells with high NKCC1 expression to an inhibitory response.⁸⁶ Preclinical studies demonstrate mixed effects, with reduction in seizure frequency and

Box 2**Principles for acute symptomatic neonatal seizure management**

Rapid and accurate electrographic seizure identification

Rapid titration of medication to abolish electrographic seizures

Early discontinuation of medication once seizures have resolved

Table 2
Pharmacologic treatment of acute symptomatic neonatal seizures

Medication	Dosage	Side Effects	Notes
<i>Levetiracetam</i>	<i>Optimal dosing not known</i> Loading dose: 40–60 mg/kg intravenously Daily dosing: 30 mg/kg/day (target levels not known) ⁹²	Mild sedation/drowsiness and irritability	Limited information regarding dosing and side effects for neonatal population
<i>Lidocaine</i>	Loading dose: 2 mg/kg over a period of 10 min, followed by a continuous infusion of 6 mg/kg/h during the first 12 h; 4 mg/kg/h for the next 12 h, and 2 mg/kg/h for the last 12 h ^a	Arrhythmia	Should only be given in the intensive care setting with continuous cardiac monitoring. In case of cardiac arrhythmia, the infusion should be discontinued immediately. Lidocaine should not be given to patients with a congenital heart disease or to neonates who have been treated with proarrhythmic drugs like phenytoin
<i>Lorazepam</i>	0.05–0.1 mg/kg intravenously	Respiratory depression, depressed level of consciousness, and hypotension	May cause myoclonus in very-low-birth-weight infants
<i>Midazolam</i>	Loading dose: 0.2 mg/kg intravenously, followed by continuous infusion (1 mcg/kg/min) increasing by 0.5–1 mcg/kg/min every 2 min to 2–5 mcg/kg/min	Respiratory depression, depressed level of consciousness, and hypotension	
<i>Phenobarbital</i>	Loading dose: 20 mg/kg intravenously, repeated once as needed Daily dosing: 5 mg/kg/d (target level 40–60 mcg/mL)	Respiratory depression, depressed level of consciousness, hypotension, and hypotonia Idiosyncratic skin rash, hepatotoxicity, and blood dyscrasias	Prolonged half-life in first week of life (43–217 hours) limits need for weaning phenobarbital in the case of short-term therapy
<i>Phenytoin and fosphenytoin</i>	Loading dose: 20 mg/kg intravenously Daily dosing: 5 mg/kg/d (target level 10–20 mcg/mL)	Infusion site reaction and arrhythmia with intravenous phenytoin Idiosyncratic skin rash, hepatotoxicity, and blood dyscrasias	Fosphenytoin has fewer cardiovascular, central nervous system, and local cutaneous side effects than phenytoin. Significant variability and changes in pharmacokinetics over the first weeks of life may lead to inconsistent drug levels

^a NB: Lower doses recommended for neonates undergoing therapeutic hypothermia.⁷⁹

duration, as well as enhanced neuroprotective efficacy when combined with phenobarbital.⁸⁷⁻⁸⁹ A single clinical study showed reduction in seizure frequency and duration following bumetanide treatment.⁹⁰ Although promising as an add-on agent for neonatal seizures, the potential for adverse effects such as ototoxicity and partial central nervous system bioavailability may ultimately limit the utility of bumetanide.⁹¹

Information about agents other than phenobarbital and phenytoin is largely derived from case series rather than randomized, blinded, clinical trials, and so the true efficacy of these medications is not known. Seizures due to acute symptomatic causes such as hypoxic-ischemic brain injury and stroke rarely persist beyond a few days of life, making any add-on agent appear more effective than the initial therapy.⁵² Furthermore, older studies do not include prolonged, cEEG monitoring, and so non-convulsive seizures, which are very common following administration of phenobarbital, may go undetected.

If the underlying cause of medically refractory seizures is unknown after initial screening laboratory tests and imaging studies, a trial of pyridoxine, pyridoxal 5'-phosphate, and folinic acid should be considered, and a screening metabolic evaluation should be performed.⁹

SUMMARY/DISCUSSION

Neonatal seizures are common and frequently reflect serious underlying brain injury. Prolonged cEEG is the gold standard for seizure monitoring; however availability remains limited at many centers. Phenobarbital, the preferred first-choice medication internationally, is effective in only 50% of cases and may be harmful, especially when used in high doses or for prolonged periods. However, there is abundant evidence from animal models to show that seizures themselves disrupt the developing brain, and so there is urgent need for research to develop safe, accurate, and widely available methods for identifying and treating electrographic seizures.

ACKNOWLEDGMENTS

Dr H.C. Glass thanks Jessica Kan for assistance with research and Dr Dawn Gano for careful review of the article.

REFERENCES

1. Annegers JF, Hauser WA, Lee JR, et al. Incidence of acute symptomatic seizures in Rochester, Minnesota, 1935-1984. *Epilepsia* 1995;36(4):327-33.
2. Jensen FE. Developmental factors regulating susceptibility to perinatal brain injury and seizures. *Curr Opin Pediatr* 2006;18(6):628-33.
3. Glass HC, Pham TN, Danielsen B, et al. Antenatal and intrapartum risk factors for seizures in term newborns: a population-based study, California 1998-2002. *J Pediatr* 2009;154(1):24-8.e1.
4. Lanska MJ, Lanska DJ, Baumann RJ, et al. A population-based study of neonatal seizures in Fayette County, Kentucky. *Neurology* 1995;45(4):724-32.
5. Saliba RM, Annegers JF, Waller DK, et al. Incidence of neonatal seizures in Harris County, Texas, 1992-1994. *Am J Epidemiol* 1999;150(7):763-9.
6. Malone A, Ryan CA, Fitzgerald A, et al. Interobserver agreement in neonatal seizure identification. *Epilepsia* 2009;50(9):2097-101.
7. Murray DM, Boylan GB, Ali I, et al. Defining the gap between electrographic seizure burden, clinical expression and staff recognition of neonatal seizures. *Arch Dis Child Fetal Neonatal Ed* 2008;93(3):F187-91.

8. Volpe JJ. Neonatal seizures. In: Volpe JJ, editor. *Neurology of the newborn*. Philadelphia: WB Saunders; 2008. p. 203–44.
9. Rahman S, Footitt EJ, Varadkar S, et al. Inborn errors of metabolism causing epilepsy. *Dev Med Child Neurol* 2013;55(1):23–36.
10. Uria-Avellanal C, Marlow N, Rennie JM. Outcome following neonatal seizures. *Semin Fetal Neonatal Med* 2013;18(4):224–32.
11. Holmes GL. The long-term effects of neonatal seizures. *Clin Perinatol* 2009;36(4):901–14, vii–viii.
12. Jensen FE. Neonatal seizures: an update on mechanisms and management. *Clin Perinatol* 2009;36(4):881–900, vii.
13. Dulac O, Mihl M, Holmes GL. Brain maturation and epilepsy. *Handb Clin Neurol* 2013;111:441–6.
14. Huttenlocher PR, de Courten C, Garey LJ, et al. Synaptogenesis in human visual cortex—evidence for synapse elimination during normal development. *Neurosci Lett* 1982;33(3):247–52.
15. Takashima S, Chan F, Becker LE, et al. Morphology of the developing visual cortex of the human infant: a quantitative and qualitative Golgi study. *J Neuropathol Exp Neurol* 1980;39(4):487–501.
16. Rakhade SN, Jensen FE. Epileptogenesis in the immature brain: emerging mechanisms. *Nat Rev Neurol* 2009;5(7):380–91.
17. Sanchez RM, Jensen FE. Maturational aspects of epilepsy mechanisms and consequences for the immature brain. *Epilepsia* 2001;42(5):577–85.
18. Dzhala VI, Staley KJ. Excitatory actions of endogenously released GABA contribute to initiation of ictal epileptiform activity in the developing hippocampus. *J Neurosci* 2003;23(5):1840–6.
19. Dzhala VI, Talos DM, Sdrulla DA, et al. NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* 2005;11(11):1205–13.
20. Khazipov R, Khalilov I, Tyzio R, et al. Developmental changes in GABAergic actions and seizure susceptibility in the rat hippocampus. *Eur J Neurosci* 2004;19(3):590–600.
21. Jiang M, Lee CL, Smith KL, et al. Spine loss and other persistent alterations of hippocampal pyramidal cell dendrites in a model of early-onset epilepsy. *J Neurosci* 1998;18(20):8356–68.
22. McCabe BK, Silveira DC, Cilio MR, et al. Reduced neurogenesis after neonatal seizures. *J Neurosci* 2001;21(6):2094–103.
23. Montgomery EM, Bardgett ME, Lall B, et al. Delayed neuronal loss after administration of intracerebroventricular kainic acid to preweanling rats. *Brain Res Dev Brain Res* 1999;112(1):107–16.
24. Lynch M, Sayin U, Bownds J, et al. Long-term consequences of early postnatal seizures on hippocampal learning and plasticity. *Eur J Neurosci* 2000;12(7):2252–64.
25. Miller SP, Weiss J, Barnwell A, et al. Seizure-associated brain injury in term newborns with perinatal asphyxia. *Neurology* 2002;58(4):542–8.
26. Glass HC, Glidden D, Jeremy RJ, et al. Clinical neonatal seizures are independently associated with outcome in infants at risk for hypoxic-ischemic brain injury. *J Pediatr* 2009;155(3):318–23.
27. Glass HC, Nash KB, Bonifacio SL, et al. Seizures and magnetic resonance imaging-detected brain injury in newborns cooled for hypoxic-ischemic encephalopathy. *J Pediatr* 2011;159(5):731–5.e1.
28. Glass HC, Bonifacio SL, Sullivan J, et al. Magnetic resonance imaging and ultrasound injury in preterm infants with seizures. *J Child Neurol* 2009;24(9):1105–11.

29. Wusthoff CJ, Drugos DJ, Gutierrez-Colina A, et al. Electrographic seizures during therapeutic hypothermia for neonatal hypoxic-ischemic encephalopathy. *J Child Neurol* 2011;26(6):724–8.
30. Clancy RR, Legido A, Lewis D. Occult neonatal seizures. *Epilepsia* 1988;29(3):256–61.
31. Bye A, Flanagan D. Electroencephalograms, clinical observations and the monitoring of neonatal seizures. *J Paediatr Child Health* 1995;31(6):503–7.
32. Shellhaas RA, Chang T, Tsuchida T, et al. The American Clinical Neurophysiology Society's guideline on continuous electroencephalography monitoring in neonates. *J Clin Neurophysiol* 2011;28(6):611–7.
33. Clancy RR. The contribution of EEG to the understanding of neonatal seizures. *Epilepsia* 1996;37(Suppl 1):S52–9.
34. Wusthoff CJ. Diagnosing neonatal seizures and status epilepticus. *J Clin Neurophysiol* 2013;30(2):115–21.
35. McCoy B, Hahn CD. Continuous EEG monitoring in the neonatal intensive care unit. *J Clin Neurophysiol* 2013;30(2):106–14.
36. Glass HC, Wusthoff CJ, Shellhaas RA. Amplitude-integrated electro-encephalography: the child neurologist's perspective. *J Child Neurol* 2013;28(10):1342–50.
37. Mastrangelo M, Fiocchi I, Fontana P, et al. Acute neonatal encephalopathy and seizures recurrence: a combined aEEG/EEG study. *Seizure* 2013;22(9):703–7.
38. Hellstrom-Westas L. Comparison between tape-recorded and amplitude-integrated EEG monitoring in sick newborn infants. *Acta Paediatr* 1992;81(10):812–9.
39. Toet MC, van der Mei W, de Vries LS, et al. Comparison between simultaneously recorded amplitude integrated electroencephalogram (cerebral function monitor) and standard electroencephalogram in neonates. *Pediatrics* 2002;109(5):772–9.
40. Shah DK, Mackay MT, Lavery S, et al. Accuracy of bedside electroencephalographic monitoring in comparison with simultaneous continuous conventional electroencephalography for seizure detection in term infants. *Pediatrics* 2008;121(6):1146–54.
41. Rennie JM, Chorley G, Boylan GB, et al. Non-expert use of the cerebral function monitor for neonatal seizure detection. *Arch Dis Child Fetal Neonatal Ed* 2004;89(1):F37–40.
42. Boylan GB, Stevenson NJ, Vanhatalo S. Monitoring neonatal seizures. *Semin Fetal Neonatal Med* 2013;18(4):202–8.
43. Temko A, Thomas E, Marnane W, et al. Performance assessment for EEG-based neonatal seizure detectors. *Clin Neurophysiol* 2011;122(3):474–82.
44. Temko A, Lightbody G, Boylan G, et al. Online EEG channel weighting for detection of seizures in the neonate. *Conf Proc IEEE Eng Med Biol Soc* 2011;2011:1447–50.
45. Temko A, Boylan G, Marnane W, et al. Robust neonatal EEG seizure detection through adaptive background modeling. *Int J Neural Syst* 2013;23(4):1350018.
46. Temko A, Stevenson N, Marnane W, et al. Inclusion of temporal priors for automated neonatal EEG classification. *J Neural Eng* 2012;9(4):046002.
47. Bonifacio SL, Miller SP. Neonatal seizures and brain imaging. *J Pediatr Neurol* 2009;7(1):61–7.
48. Booth D, Evans DJ. Anticonvulsants for neonates with seizures. *Cochrane Database Syst Rev* 2004;(4):CD004218.

49. Slaughter LA, Patel AD, Slaughter JL. Pharmacological treatment of neonatal seizures: a systematic review. *J Child Neurol* 2013;28(3):351–64.
50. Bittigau P, Siffringer M, Genz K, et al. Antiepileptic drugs and apoptotic neurodegeneration in the developing brain. *Proc Natl Acad Sci U S A* 2002;99(23):15089–94.
51. Painter MJ, Scher MS, Stein AD, et al. Phenobarbital compared with phenytoin for the treatment of neonatal seizures. *N Engl J Med* 1999;341(7):485–9.
52. Lynch NE, Stevenson NJ, Livingstone V, et al. The temporal evolution of electrographic seizure burden in neonatal hypoxic ischemic encephalopathy. *Epilepsia* 2012;53(3):549–57.
53. Grimshaw JM, Russell IT. Effect of clinical guidelines on medical practice: a systematic review of rigorous evaluations. *Lancet* 1993;342(8883):1317–22.
54. Guillet R, Kwon JM. Prophylactic phenobarbital administration after resolution of neonatal seizures: survey of current practice. *Pediatrics* 2008;122(4):731–5.
55. Hellstrom-Westas L, Blennow G, Lindroth M, et al. Low risk of seizure recurrence after early withdrawal of antiepileptic treatment in the neonatal period. *Arch Dis Child Fetal Neonatal Ed* 1995;72(2):F97–101.
56. Guillet R, Kwon J. Seizure recurrence and developmental disabilities after neonatal seizures: outcomes are unrelated to use of phenobarbital prophylaxis. *J Child Neurol* 2007;22(4):389–95.
57. Bartha AI, Shen J, Katz KH, et al. Neonatal seizures: multicenter variability in current treatment practices. *Pediatr Neurol* 2007;37(2):85–90.
58. Wheless JW, Clarke DF, Arzimanoglou A, et al. Treatment of pediatric epilepsy: European expert opinion, 2007. *Epileptic Disord* 2007;9(4):353–412.
59. Bassan H, Bentol Y, Shany E, et al. Neonatal seizures: dilemmas in workup and management. *Pediatr Neurol* 2008;38(6):415–21.
60. Boylan GB, Rennie JM, Chorley G, et al. Second-line anticonvulsant treatment of neonatal seizures: a video-EEG monitoring study. *Neurology* 2004;62(3):486–8.
61. Glass HC, Kan J, Bonifacio SL, et al. Neonatal seizures: treatment practices among term and preterm infants. *Pediatr Neurol* 2012;46(2):111–5.
62. Silverstein FS, Ferriero DM. Off-label use of antiepileptic drugs for the treatment of neonatal seizures. *Pediatr Neurol* 2008;39(2):77–9.
63. Mbizvo GK, Dixon P, Hutton JL, et al. Levetiracetam add-on for drug-resistant focal epilepsy: an updated Cochrane Review. *Cochrane Database Syst Rev* 2012;(9):CD001901.
64. Kim JS, Kondratyev A, Tomita Y, et al. Neurodevelopmental impact of antiepileptic drugs and seizures in the immature brain. *Epilepsia* 2007;48(Suppl 5):19–26.
65. Manthey D, Asimiadou S, Stefovská V, et al. Sulthiame but not levetiracetam exerts neurotoxic effect in the developing rat brain. *Exp Neurol* 2005;193(2):497–503.
66. Talos DM, Chang M, Kosaras B, et al. Antiepileptic effects of levetiracetam in a rodent neonatal seizure model. *Pediatr Res* 2013;73(1):24–30.
67. Kilicdag H, Daglioglu K, Erdogan S, et al. The effect of levetiracetam on neuronal apoptosis in neonatal rat model of hypoxic ischemic brain injury. *Early Hum Dev* 2013;89(5):355–60.
68. Tulloch JK, Carr RR, Ensom MH. A systematic review of the pharmacokinetics of antiepileptic drugs in neonates with refractory seizures. *J Pediatr Pharmacol Ther* 2012;17(1):31–44.
69. Merhar SL, Schibler KR, Sherwin CM, et al. Pharmacokinetics of levetiracetam in neonates with seizures. *J Pediatr* 2011;159(1):152–4.e3.

70. Sharpe CM, Capparelli EV, Mower A, et al. A seven-day study of the pharmacokinetics of intravenous levetiracetam in neonates: marked changes in pharmacokinetics occur during the first week of life. *Pediatr Res* 2012;72(1):43–9.
71. Abend NS, Gutierrez-Colina AM, Monk HM, et al. Levetiracetam for treatment of neonatal seizures. *J Child Neurol* 2011;26(4):465–70.
72. Khan O, Chang E, Cipriani C, et al. Use of intravenous levetiracetam for management of acute seizures in neonates. *Pediatr Neurol* 2011;44(4):265–9.
73. Furwentsches A, Bussmann C, Ramantani G, et al. Levetiracetam in the treatment of neonatal seizures: a pilot study. *Seizure* 2010;19(3):185–9.
74. Sheth RD, Buckley DJ, Gutierrez AR, et al. Midazolam in the treatment of refractory neonatal seizures. *Clin Neuropharmacol* 1996;19(2):165–70.
75. Hu KC, Chiu NC, Ho CS, et al. Continuous midazolam infusion in the treatment of uncontrollable neonatal seizures. *Acta Paediatr Taiwan* 2003;44(5):279–81.
76. Castro Conde JR, Hernandez Borges AA, Domenech Martinez E, et al. Midazolam in neonatal seizures with no response to phenobarbital. *Neurology* 2005;64(5):876–9.
77. Sirsi D, Nangia S, LaMothe J, et al. Successful management of refractory neonatal seizures with midazolam. *J Child Neurol* 2008;23(6):706–9.
78. Lundqvist M, Agren J, Hellstrom-Westas L, et al. Efficacy and safety of lidocaine for treatment of neonatal seizures. *Acta Paediatr* 2013;102(9):863–7.
79. van den Broek MP, Rademaker CM, Straaten HL, et al. Anticonvulsant treatment of asphyxiated newborns under hypothermia with lidocaine: efficacy, safety and dosing. *Arch Dis Child Fetal Neonatal Ed* 2013;98(4):F341–5.
80. van den Broek MP, Huitema AD, van Hasselt JG, et al. Lidocaine (lignocaine) dosing regimen based upon a population pharmacokinetic model for preterm and term neonates with seizures. *Clin Pharmacokinet* 2011;50(7):461–9.
81. Malingre MM, Van Rooij LG, Rademaker CM, et al. Development of an optimal lidocaine infusion strategy for neonatal seizures. *Eur J Pediatr* 2006;165(9):598–604.
82. Cha BH, Silveira DC, Liu X, et al. Effect of topiramate following recurrent and prolonged seizures during early development. *Epilepsy Res* 2002;51(3):217–32.
83. Liu Y, Barks JD, Xu G, et al. Topiramate extends the therapeutic window for hypothermia-mediated neuroprotection after stroke in neonatal rats. *Stroke* 2004;35(6):1460–5.
84. Clark AM, Kriel RL, Leppik IE, et al. Intravenous topiramate: comparison of pharmacokinetics and safety with the oral formulation in healthy volunteers. *Epilepsia* 2013;54(6):1099–105.
85. Clark AM, Kriel RL, Leppik IE, et al. Intravenous topiramate: safety and pharmacokinetics following a single dose in patients with epilepsy or migraines taking oral topiramate. *Epilepsia* 2013;54(6):1106–11.
86. Khanna A, Walcott BP, Kahle KT. Limitations of current GABA agonists in neonatal seizures: toward GABA modulation via the targeting of neuronal Cl⁽⁻⁾ transport. *Front Neurol* 2013;4:78.
87. Dzhala VI, Brumback AC, Staley KJ. Bumetanide enhances phenobarbital efficacy in a neonatal seizure model. *Ann Neurol* 2008;63(2):222–35.
88. Cleary RT, Sun H, Huynh T, et al. Bumetanide enhances phenobarbital efficacy in a rat model of hypoxic neonatal seizures. *PLoS One* 2013;8(3):e57148.
89. Liu Y, Shangguan Y, Barks JD, et al. Bumetanide augments the neuroprotective efficacy of phenobarbital plus hypothermia in a neonatal hypoxia-ischemia model. *Pediatr Res* 2012;71(5):559–65.

90. Kahle KT, Barnett SM, Sassower KC, et al. Decreased seizure activity in a human neonate treated with bumetanide, an inhibitor of the Na(+)–K(+)–2Cl(–) co-transporter NKCC1. *J Child Neurol* 2009;24(5):572–6.
91. Chabwine JN, Vanden Eijnden S. A claim for caution in the use of promising bumetanide to treat neonatal seizures. *J Child Neurol* 2011;26(5):657–8 [author reply: 658–9].
92. ClinicalTrials.gov. Identifier: NCT01720667.